



July 15, 2020

Dear Parent or Guardian:

As school nurse at the Waunakee Community Intermediate School, I am writing to inform you about the dangers of meningococcal disease, commonly known as bacterial meningitis, a rare but potentially fatal infection that can occur among teenagers and college students. State Statute requires each school district to provide parents/guardians of students enrolled in grade 6 with information about meningococcal disease. While meningococcal disease is rare and difficult to contract, it is very serious. There is now a vaccine that may help to prevent this infection.

Meningococcal bacteria can potentially be transmitted through close contact with an infected person through direct contact with respiratory and/or oral secretions from an infected person (for example, through sharing drinking containers or kissing). Teenagers and college students are at increased risk for meningococcal disease compared to the general population, accounting for nearly 30 percent of all U.S. cases every year. Meningococcal disease can be misdiagnosed as something less serious, because early symptoms like high fever, severe headache, nausea, vomiting and stiff neck, are similar to those of common viral illnesses. The disease can progress rapidly and can cause death or permanent disability within 48 hours of initial symptoms.

Up to 83 percent of all cases among teens and college students may potentially be prevented through immunization, the most effective way to prevent this disease. A meningococcal vaccine is available that protects against four out of five strains of bacterium that cause meningococcal disease in the U.S.

The Centers for Disease Control and Prevention (CDC) and other leading medical organizations recommends that all 11-12 years olds should be vaccinated with meningococcal conjugate vaccine (MCV4). A booster shot is recommended for teens at age 16 to continue providing protection when their risk for meningococcal disease is highest. Teens who received MCV4 for the first time at age 13 through 15 years will need a one-time booster dose at 16 through 18 years of age. If a teenager missed getting the vaccine altogether, they should ask the doctor about getting it now, especially if they are about to move into a college dorm or military barracks.

For more information about meningococcal disease and immunization, please feel free to contact me directly at 608-849-2176 Ext. 2 or visit the following website, https://dpi.wi.gov/sites/default/files/imce/sspw/doc/snmening_doc to learn more about meningococcal disease, vaccine information, and public health resources visit the following web sites. I also encourage you to discuss this with your child's primary care physician.

Sincerely,

Jennifer Sydow and Sara Anderson-Kim
School Nurse

Chris Mand
Associate Director of Student Services

Kurt Eley
Director of Student Services

Meningococcal Disease: Protect Your Child

Public health authorities recommend that teenagers and college-bound students be immunized against a potentially fatal bacterial infection called meningococcal disease, a type of meningitis.

Meningococcal disease is a rare but potentially fatal bacterial infection that can cause severe swelling of the brain and spinal cord (meningitis) or a serious blood infection (meningococemia). Meningococcal disease strikes up to 3,000 Americans each year; nearly 30 percent of these cases are among teenagers and college students.

Up to 83 percent of all cases among teens and college students may potentially be prevented through immunization, the most effective way to prevent this disease. A meningococcal vaccine is available that protects against four out of five strains of bacterium that cause meningococcal disease in the U.S.

The Centers for Disease Control and Prevention (CDC) and other leading medical organizations recommends that all 11-12 years olds should be vaccinated with meningococcal conjugate vaccine (MCV4). A booster shot is recommended for teens at age 16 to continue providing protection when their risk for meningococcal disease is highest. Teens who received MCV4 for the first time at age 13 through 15 years will need a one-time booster dose at 16 through 18 years of age. If a teenager missed getting the vaccine altogether, they should ask the doctor about getting it now, especially if they are about to move into a college dorm or military barracks.

About Meningococcal Disease

Meningococcal disease is often misdiagnosed as something less serious because early symptoms are similar to common viral illnesses. Symptoms of meningococcal disease may include high fever, severe headache, stiff neck, nausea, vomiting, sensitivity to light, confusion, exhaustion and/or a rash.

Teenagers and college students are at increased risk for meningococcal disease compared to the general population, accounting for nearly 30 percent of all U.S. cases every year. Meningococcal disease can be misdiagnosed as something less serious, because early symptoms like high fever, severe headache, nausea, vomiting and stiff neck, are similar to those of common viral illnesses. The disease can progress rapidly and can cause death or permanent disability within 48 hours of initial symptoms.

Meningococcal disease is spread through direct contact with respiratory and/or oral secretions from infected persons (for example, kissing or sharing drinking containers). It can develop and spread quickly throughout the body, so early diagnosis and treatment are very important. Even with immediate treatment, the disease can kill an otherwise healthy young person within hours of first symptoms. Of those who survive, up to 20 percent may endure permanent disabilities, including brain damage, deafness and limb amputations.

Lifestyle factors common among teenagers, college students and military personnel are believed to put them at increased risk of contracting meningococcal disease. These lifestyle factors include crowded living situations (for example, dormitories, sleep-away camps), active or passive smoking and irregular sleeping habits. Teens should avoid sharing eating utensils and drinking out of the same container, since infections may spread through this type of close contact.

To learn more about meningococcal disease, vaccine information, and public health resources visit the following web sites.

Center for Disease Control and Prevention meningococcal meningitis information.

- General information regarding [meningitis disease](#)
- A [Meningitis Fact Sheet](#) (This is a pdf document that cannot be altered, but may be copied and distributed.)
- American Committee of Immunization Practice recommendations for Prevention and Control of Meningitis (2005)
<http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5407a1.htm>
- Updated recommendations for the use of Meningococcal Conjugate Vaccines (2010) is available at:
http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6003a3.htm?s_cid=mm6003a3_e

National Association of School Nurses – Voices of Meningitis

(<http://www.nasn.org/ToolsResources/Immunizations/VoicesofMeningitisChallenge>)

- General parent letter [[English](#)] [[Spanish](#)]
- Fact Sheet [[English](#)] [[Spanish](#)]
- Vaccine Information Statement ([VIS](#))
- [Presentation to parents](#)

A list of local [Wisconsin public health departments and contact information](#)

- [Meningitis Foundation of America](#)
- [National Meningitis Association](#)
- [American Academy of Family Physicians](#)
- [American Academy of Pediatrics](#)